

Research

Open Access

Gastric intramucosal pH is stable during titration of positive end-expiratory pressure to improve oxygenation in acute respiratory distress syndrome

Ibrahim Ozkan Akinci¹, Nahit Çakar², Gökhan Mehmet Mutlu³, Simru Tugrul¹,
Perihan Ergin Ozcan¹, Musa Gitmez¹, Figen Esen² and Lutfi Telci²

¹Attendings of Anesthesiology and Intensive Care, Department of Anesthesiology and Intensive Care, Istanbul Medical Faculty, Capa Klinikleri, Istanbul, Turkey

²Professor of Anesthesiology and Intensive Care, Department of Anesthesiology and Intensive Care, Istanbul Medical Faculty, Capa Klinikleri, Istanbul, Turkey

³Assistant Professor of Medicine, Pulmonary and Critical Care Medicine, Evanston Northwestern Healthcare, Evanston, and Northwestern University, Illinois, USA

Correspondence: I Ozkan Akinci, iozkana@yahoo.com

Received: 12 February 2003

Revisions requested: 20 February 2003

Revisions received: 24 February 2003

Accepted: 24 February 2003

Published: 12 March 2003

Critical Care 2003, **7**:R17-R23 (DOI 10.1186/cc2172)

This article is online at <http://ccforum.com/content/7/3/R17>

© 2003 Akinci *et al.*, licensee BioMed Central Ltd (Print ISSN 1364-8535; Online ISSN 1466-609X). This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background Optimal positive end-expiratory pressure (PEEP) is an important component of adequate mechanical ventilation in acute lung injury and acute respiratory distress syndrome (ARDS). In the present study we tested the effect on gastric intramucosal pH of incremental increases in PEEP level (i.e. PEEP titration) to improve oxygenation in ARDS. Seventeen consecutive patients with ARDS, as defined by consensus criteria, were included in this clinical, prospective study. All patients were haemodynamically stable and were not receiving vasopressors. From an initial level of 5 cmH₂O, PEEP was titrated at 2 cmH₂O increments until the partial arterial oxygen tension was 300 mmHg or greater, peak airway pressure was 45 cmH₂O or greater, or mean arterial blood pressure decreased by 20% or more of the baseline value. Optimal PEEP was defined as the level of PEEP that achieved the best oxygenation. The maximum PEEP was the highest PEEP level reached during titration in each patient.

Results Gastric mucosal pH was measured using gastric tonometry at all levels of PEEP. The thermodilution technique was used for measurement of cardiac index. Gastric mucosal pH was similar at baseline and at optimal PEEP levels, but it was slightly reduced at maximum PEEP. Cardiac index and oxygen delivery remained stable at all PEEP levels.

Conclusion Incremental titration of PEEP based on improvement in oxygenation does not decrease gastric intramucosal perfusion when cardiac output is preserved in patients with ARDS.

Keywords acute lung injury, acute respiratory distress syndrome, mechanical ventilation, positive end-expiratory pressure, splanchnic perfusion

Introduction

Positive end-expiratory pressure (PEEP) is an important component of the ventilatory management of acute lung injury

(ALI) and acute respiratory distress syndrome (ARDS). PEEP improves oxygenation by redistributing the alveolar fluid and restores functional residual capacity by keeping the alveoli

ALI = acute lung injury; ARDS = acute respiratory distress syndrome; DO₂ = oxygen delivery; MAP = mean arterial pressure; Pao₂ = partial arterial oxygen tension; P_(t-a)CO₂ = gap between partial tissue and arterial carbon dioxide tension; PEEP = positive end-expiratory pressure; pH_i = gastric mucosal pH.

open. However, PEEP can be detrimental because it may, particularly at high levels, decrease cardiac output by decreasing the venous return as a result of diminished pressure gradient between the systemic veins and right atrium [1], and consequently it may lead to hypoperfusion of vital organs. Ultimately, despite improving arterial oxygen content, PEEP may decrease oxygen delivery to various organs, among which the splanchnic vascular bed appears to be particularly at risk because of its predisposing features and the influence of PEEP on regional blood flow distribution.

Maintenance of splanchnic blood flow is important because splanchnic hypoperfusion may play a critical role in the pathogenesis of multiorgan dysfunction syndrome [2,3]. Mechanical ventilation has been suggested to potentiate the adverse effects of underlying critical illness on splanchnic vasculature and contribute to the development of multiorgan dysfunction syndrome, particularly when 'injurious' ventilatory strategies that produce high end-inspiratory lung volumes are employed [3]. Experimental studies suggested that mechanical ventilation with considerably high levels of PEEP can lead to splanchnic hypoperfusion and marked reduction in hepatic blood flow [4–6]. Furthermore, PEEP may decrease splanchnic blood flow in patients with no underlying lung disease [7,8]. Most available evidence regarding the effects of PEEP from animal studies has been extrapolated to humans based on the assumption that the effects of mechanical ventilation on humans and animals are similar. However, a recent study conducted in humans explored the effect of PEEP in patients with ALI [9] and did not find a consistent effect on splanchnic blood flow.

Because of the difficulties associated with measurement of pressure–volume curves, incremental titration of PEEP in an attempt to find the 'best' PEEP, based on improvement in oxygenation, is common practice in the management of hypoxaemic respiratory failure. However, it is unknown whether this strategy has an adverse effect on splanchnic perfusion. The aim of the present study was to investigate the impact of PEEP titration (based on improvement in oxygenation) on gastric mucosal perfusion in patients with ARDS, as assessed by measurement of gastric mucosal pH (pH_i).

Method

Patients

The study protocol was approved by the Institutional Ethics Committee of Istanbul University Hospital. Written informed consent was obtained from each patient or the patient's next of kin. We consecutively enrolled 17 patients with ARDS admitted to the multidisciplinary intensive care unit at Istanbul University Hospital. The criteria for eligibility were a diagnosis of ARDS (based on a consensus report [10]), age older than 18 years and mean arterial pressure (MAP) greater than 60 mmHg with no haemodynamic support. All patients were enrolled within the first 24 hours following the diagnosis of ARDS. Patients with known cardiac dysfunction or pre-existing liver disease were not included in the trial.

Protocol

All patients were ventilated using a Servo 300 Siemens ventilator (Siemens Elema, Uppsala, Sweden) using the pressure-regulated volume control mode with a tidal volume of 8–10 ml/kg (based on ideal body weight), frequency of 12 breaths/min, fraction of inspired oxygen of 1.0, and inspiratory:expiratory ratio of 1:2. Patients were sedated with midazolam (Dormicum; Hoffmann LaRoche, Basel, Switzerland) at 4 mg/hour and paralyzed with 0.1 mg/kg vecuronium (Norcuron; Organon, Oss, The Netherlands) infusion during the study. In addition to employing a radial arterial catheter for blood pressure measurement, a pulmonary artery catheter (Abbot Labs, North Chicago, IL, USA) was placed in all patients for haemodynamic monitoring. No patients received any therapeutic intervention to improve haemodynamics (i.e. fluid resuscitation or catecholamine infusion) throughout the study.

Baseline PEEP ($PEEP_{baseline}$) was set at 5 cmH₂O and titrated at 2 cmH₂O increments until the partial arterial oxygen tension (PaO_2) reached at least 300 mmHg, peak airway pressure was 45 cmH₂O or greater, or MAP dropped by 20% or more from the baseline value. Criteria for overinflation of lung (and therefore for discontinuation of further titration of PEEP) were reduction in PaO_2 of 10% or more and an increase in arterial carbon dioxide tension of 10% or more. Optimal PEEP ($PEEP_{opt}$) was defined as the PEEP that achieved the best oxygenation, whereas maximum PEEP ($PEEP_{max}$) was the greatest level of PEEP achieved during titration in each patient.

A nasogastric catheter (TRIP Catheter; Tonometrics Division, Instrumentarium Corp., Helsinki, Finland) was inserted into the stomach to measure pH_i . Correct placement of the TRIP catheter was confirmed by radiography. Enteral nutrition was withheld throughout the study, and all patients received ranitidine 50 mg intravenously. In order to allow for equilibration, pH_i was measured 45 min after injection of 2.5 ml isotonic saline into the semipermeable balloon of the TRIP catheter. Partial pressure of carbon dioxide in saline solution and bicarbonate level in arterial blood were measured simultaneously using a blood gas analyzer (ABL-500; Radiometer, Copenhagen, Denmark) immediately after sampling [11] and were corrected for the equilibration time [12]. The pH_i was calculated using the Henderson–Hasselbach equation.

All measurements, including respiratory, haemodynamic parameters, arterial and mixed venous blood gas analyses, and gastric pH_i , were taken at baseline and following ventilation for 45 min at each level of PEEP. Haemodynamic parameters were monitored continuously using an Horizon XL monitor (Mennen Medical Inc., New York, NY, USA). Cardiac output was measured in triplicate by thermodilution technique using 10 ml saline solution at room temperature. Cardiac index, shunt fraction, oxygen delivery (DO_2) and oxygen consumption were calculated at baseline and at all PEEP levels.

Table 1**Characteristics of the 17 patients studied**

Patient number	Diagnosis at admission	Age (years)	Sex	APACHE II score	SOFA score	Additional organ failure
1	Multiple trauma	74	M	15	5	R, N
2	Hepatic coma	26	F	23	10	R, H, N, L
3	Cerebral ischaemia	70	M	22	6	H, N
4	Sepsis	55	M	21	6	H, L
5	Intracranial haemorrhage	58	M	15	7	H, N
6	Acute pancreatitis	50	M	23	9	R, H, L
7	Multiple trauma	24	M	19	6	R, H
8	Pneumonia	18	F	12	5	R
9	Postoperative sepsis	28	F	21	7	R, H, N, L
10	Acute pancreatitis	74	M	23	7	R, H, L
11	Tetanus	62	M	22	4	R, H
12	Pneumonia	62	M	21	3	H
13	Bronchopneumonia	37	F	16	4	R
14	Multiple trauma	40	M	21	5	R, H
15	Postoperative scoliosis	19	F	24	7	R, H, L
16	Aortoduodenal fistula	69	M	17	6	R, H, L
17	Intra-abdominal sepsis	37	F	20	7	R, H, L

APACHE, Acute Physiology and Chronic Health Evaluation; F, female; H, haematological system; L, hepatic system; M, male; N, neurological system; R, renal system; SOFA, Sequential Organ Failure Assessment.

Statistical analysis

Paired analysis of variance tests were used to analyze the differences between measurements. $P < 0.05$ was considered statistically significant. All values are presented as mean \pm standard deviation.

Results

A total of 17 patients were enrolled in the present study (11 male and 6 female). The characteristics of the individual patients are shown in Table 1. The mean age of the study population was 47.2 ± 19.8 , the mean Acute Physiology and Chronic Health Evaluation II score was 19.7 ± 3.5 , and the mean Sequential Organ Failure Assessment score was 6.3 ± 1.8 . By titrating PEEP, we were able to achieve a mean PEEP_{opt} of 10.4 ± 3.9 cmH₂O and a PEEP_{max} of 13.3 ± 2.9 cmH₂O ($P = 0.0001$). The highest PEEP value applied was 17 cmH₂O. Static compliance improved slightly at PEEP_{opt}, but this did not achieve statistical significance ($P = 0.84$; Table 2). Changes in peak airway and mean airway pressures at PEEP_{baseline}, PEEP_{opt} and PEEP_{max} were statistically significant ($P < 0.001$; Table 2). Reasons for stopping the titration of PEEP were reduction in PaO₂ (from 20% to 40%; $n = 6$), reduction in MAP (from 25% to 60%; $n = 4$), adequate oxygenation (PaO₂ 350–450 mmHg; $n = 4$) and excessive peak upper airway pressure ($n = 3$).

Although PEEP significantly improved shunt fraction, and consequently PaO₂, its greater effect on cardiac output led to a reduction in DO₂ both at PEEP_{opt} and PEEP_{max}. However, none of the changes in haemodynamic parameters, including those in central venous pressure, pulmonary artery occlusion pressure, cardiac output, cardiac index and DO₂, achieved statistical significance (Table 2 and Fig. 1). PaO₂ values remained stable at each level of PEEP. The mean pH_i was 7.31 ± 0.13 at baseline and 7.32 ± 0.12 at PEEP_{opt}; it decreased to 7.29 ± 0.12 at PEEP_{max}, but this reduction was not statistically significant ($P = 0.84$). Similar to pH_i, alterations in the gap between partial tissue and arterial carbon dioxide tension ($P_{(t-a)}\text{CO}_2$) were not significant ($P = 0.353$).

Although the increase in PEEP had no impact on the group as a whole, changes in pH_i and $P_{(t-a)}\text{CO}_2$ during PEEP titration differed between individual patients (Table 3). The pH_i decreased in eight patients (47%), it increased in five (29.4%) and it was unchanged in four (23.5%) at PEEP_{opt} as compared with PEEP_{baseline}. The pH_i at PEEP_{max} was lower in 12 (70.6%) and higher in five (29.4%) patients as compared with baseline values. The $P_{(t-a)}\text{CO}_2$ values increased in nine (52.3%) patients at PEEP_{opt} and in 10 (58.3%) patients at PEEP_{max} as compared with PEEP_{baseline} (Table 3). However, there were no statistically significant differences in $P_{(t-a)}\text{CO}_2$

Table 2**Parameters measured during titration of positive end-expiratory pressure**

Parameter	PEEP _{baseline}	PEEP _{opt}	PEEP _{max}	P value
PEEP (cmH ₂ O)	5	10.4 ± 3.9	13.3 ± 2.9	0.0001
P _{peak} (cmH ₂ O)	27.2 ± 5	31.5 ± 6.2	35 ± 5	0.0001
P _{mean} (cmH ₂ O)	11.4 ± 1.9	15.9 ± 4.9	19.3 ± 2.8	0.0001
PaO ₂ (mmHg)	136.6 ± 48.7	231 ± 86.1	226 ± 99.8	0.001
Paco ₂ (mmHg)	38.4 ± 7.02	37 ± 7.63	37.4 ± 8.19	0.70
pH _i	7.31 ± 0.13	7.32 ± 0.12	7.30 ± 0.12	0.84
P _(t-a) CO ₂ (mmHg)	3.74 ± 8.31	5.27 ± 5.49	7.42 ± 7.39	0.353
CI (l/min per m ²)	4.03 ± 1.47	3.72 ± 1.4	3.62 ± 1.21	0.79
CO (l/min)	7.02 ± 2.46	6.61 ± 2.41	6.5 ± 2.15	0.13
MAP (mmHg)	89 ± 17.7	88.4 ± 15.3	83 ± 15.9	0.49
CVP (mmHg)	13 ± 2.9	12 ± 3.4	11.5 ± 2.8	0.45
PCWP (mmHg)	15 ± 3.3	12.7 ± 3.07	12.5 ± 3.1	0.10
DO ₂ (ml/min per m ²)	689 ± 232.9	659.9 ± 221.7	638.5 ± 197.4	0.14
VO ₂ (ml/min per m ²)	244.2 ± 73.4	233 ± 42.08	251.2 ± 49.7	0.84
Qs/Qt (%)	34.41 ± 6.23	23.1 ± 9.18	27.4 ± 6.65	0.03
O ₂ ext (%)	21.88 ± 4.65	27.03 ± 3.22	28.51 ± 8.53	0.57
C _{st} (ml/cmH ₂ O)	32.3 ± 9.7	33 ± 8.35	32.9 ± 8.5	0.84


CI, cardiac index; CO, cardiac output; C_{st}, static compliance; CVP, central venous pressure; DO₂, oxygen delivery; MAP, mean arterial pressure; O₂ext, oxygen extraction ratio; PaO₂, partial arterial oxygen tension; PCWP, pulmonary capillary wedge pressure; PEEP, positive end-expiratory pressure; pH_i, gastric mucosal pH; P_{mean}, mean airway pressure; P_(t-a)CO₂, gap between partial tissue and arterial carbon dioxide tension; P_{peak}, peak airway pressure; Qs/Qt, shunt fraction; VO₂, oxygen consumption.

values between PEEP_{baseline}, PEEP_{opt} and PEEP_{max} ($P=0.353$; Table 2). Interestingly, DO₂ in those patients who exhibited a rise in pH_i did not increase. Rather, DO₂ in these patients also decreased (although this was not statistically significant) at PEEP_{opt} and PEEP_{max} to a degree similar to that in patients who exhibited a drop in pH_i.

Discussion

The results of the present study indicate that incremental increases in PEEP do not impact on splanchnic perfusion, as assessed by gastric tonometry, when cardiac output (and consequently DO₂) is maintained.

In animals, PEEP decreases hepatosplanchnic perfusion in a dose-dependent manner, with a limited effect at PEEP levels of less than 10 cmH₂O [2,4,5]. Alterations in splanchnic blood flow attributed to PEEP occur in parallel to those in cardiac output and consequently can be reversed with restoration of blood pressure [4,13]. Despite experimental evidence, concerns regarding the effects of PEEP on splanchnic perfusion remain theoretical because large studies in humans are lacking. Similarly, in humans without ALI or ARDS, PEEP reduces splanchnic oxygenation and this is accompanied by decreases in cardiac output, albeit with

Figure 1

Cardiac output changes at baseline positive end-expiratory pressure (PEEP_{baseline}: 5 cmH₂O), PEEP_{opt} and PEEP_{max}

no change in lactate levels [14]. Recently, Kiefer and colleagues [9] reported no change in splanchnic perfusion when PEEP was titrated on the linear portion of the pressure–volume curve in patients with ALI [9].

Table 3

Levels of positive end-expiratory pressure achieved and corresponding levels of gastric mucosal pH, and partial tissue and arterial carbon dioxide tension gap

Patient number	PEEP _{opt} (cmH ₂ O)	PEEP _{max} (cmH ₂ O)	pH _i			P _(t-a) CO ₂		
			At PEEP _{baseline}	At PEEP _{opt}	At PEEP _{max}	At PEEP _{baseline}	At PEEP _{opt}	At PEEP _{max}
1	7	9	7.46	7.4	7.4	-1.7	-1	-3.5
2	13	15	7.45	7.43	7.33	-1.7	6.7	17.5
3	17	17	7.26	7.28	7.28	3	-2	-2
4	5	15	7.19	7.19	7.16	10.2	8	13
5	7	11	7.45	7.37	7.32	-3.4	3.4	11
6	15	15	7.23	7.05	7.05	-0.7	11.6	12.6
7	7	15	7.24	7.26	7.23	1.5	0.9	16.1
8	9	13	7.18	7.18	7.24	11.2	14	10.8
9	17	17	7.45	7.44	7.44	-1	10	5
10	15	17	7.15	7.23	7.2	4.6	10.5	13.2
11	11	15	7.45	7.42	7.29	-4.7	5	8.9
12	13	15	7.50	7.43	7.46	-0.3	8	-3.6
13	13	13	7.27	7.46	7.46	-3.5	1	7.3
14	5	9	7.36	7.36	7.27	3.4	3.8	7.4
15	7	9	7.38	7.37	7.37	2.1	13	16
16	5	13	7.14	7.14	7.08	7	-3.8	-2.6
17	9	9	7.15	7.46	7.46	26.6	0.5	-1

Positive end-expiratory pressure (PEEP) at baseline was 5 cmH₂O. pH_i, gastric mucosal pH; P_(t-a)CO₂, gap between partial tissue and arterial carbon dioxide tension.

The results presented here, which demonstrate a lack of impact on splanchnic blood flow when PEEP is not accompanied by decreased cardiac output, corroborate those from animal studies [4,13] and from the recent human study conducted by Kiefer and coworkers [9]. The lack of change in pH_i at PEEP_{opt} (11 cmH₂O) is in agreement with our current understanding that PEEP at 10 cmH₂O has a limited effect on splanchnic blood flow. Furthermore, the presence of ARDS limited the relative impact of increased thoracic pressure on the cardiovascular system.

Perhaps more important, these observations were valid for a wide range of PEEP levels, from 5 cmH₂O to as high as 17 cmH₂O. We ascribed the lack of significant changes in cardiac output and DO₂ in the patients to adequate volume status and preload. Relative hypovolaemia appears to be the most likely explanation for the reductions in cardiac output and splanchnic blood flow observed in animal studies. Gastric pH_i, and consequently splanchnic blood flow, remained stable at PEEP_{opt} and PEEP_{max} when cardiac output and DO₂ remained relatively unchanged. Preservation of splanchnic blood flow at PEEP_{opt} and PEEP_{max} was attributed to an increase in oxygen extraction ratio that was sufficient to compensate for the small,

insignificant drop in cardiac output and DO₂ that occurred during PEEP titration [15].

It is also noteworthy that there may be individual variations in pH_i in response to PEEP. Although differences in pH_i response among individuals cannot be explained on the basis of changes in DO₂, they may be attributed to differences in the relative impact of underlying critical illness on splanchnic perfusion and variations in splanchnic vascular response (i.e. severity and/or duration of vasoconstriction, extraction ratio) to small changes in DO₂ among individuals.

Because of concerns about the reliability of pH_i for assessing mucosal perfusion, we also calculated the P_(t-a)CO₂ because it has been proposed to be a better parameter than pH_i [16]. The pH_i level can sometimes be misleading, particularly in situations in which gastric tissue and arterial bicarbonate levels are not equal. In addition, unlike pH_i, which can change with the degree of alveolar ventilation, P_(t-a)CO₂ remains a reliable parameter because both components (i.e. partial arterial and tissue carbon dioxide tension) are similarly influenced by changes in alveolar ventilation, unless they are associated with alterations in cardiac output [17]. In the present study,

changes in $P_{(t-a)}\text{CO}_2$ were not statistically significant and correlated with changes in pH_i . Consequently, we used pH_i values in our discussion because we believe that pH_i reliably reflects the accurate tissue pH in patients.

Our results corroborate those from the only other study that evaluated the impact of PEEP on splanchnic perfusion in patients with ALI. Similar to Kiefer and colleagues [9], we found no change either in pH_i or $P_{(t-a)}\text{CO}_2$ during PEEP titration. However, there were several differences between two studies. Whereas Kiefer and colleagues used pressure-volume curves for PEEP titration, we titrated PEEP on the basis of improvement in oxygenation, which is a commonly used method in clinical practice because determination of pressure-volume curves can sometimes be cumbersome. Furthermore, the present study was larger and we included patients with more severe disease (ratio of fractional inspired oxygen to PaO_2 : 139 in the present study versus 168 in that conducted by Kiefer and coworkers).

However, the present study has several limitations. The first and perhaps most important limitation of the study is the liberal titration of PEEP in order to determine its impact on pH_i , as described under Method (see above). We acknowledge that in day-to-day clinical practice, some of the patients would not have been managed with such aggressive titration of PEEP and therefore would not have received the levels of PEEP achieved in the study, rendering the clinical implications of these observations quite limited. Second, we did not directly measure splanchnic perfusion but assessed it indirectly by monitoring pH_i using gastric tonometry. Although the diagnostic value of gastric tonometry has been questioned because of some methodological problems, we believe that we minimized most of these limitations and improved the reproducibility of our measurements by immediate analysis of samples, use of H_2 blockers [18] and lack of enteral nutrition [19], rendering it possible to use gastric pH_i to evaluate splanchnic perfusion. Third, PEEP_{opt} in the study (approximately 11 cmH_2O) was lower than levels reported in other ARDS studies [20]. Higher tidal volume (10 ml/kg) leading to higher mean airway pressure, the termination criteria used in our study, and the differences in titration technique (based on oxygenation versus pressure-volume curve) may account for this difference. Finally, pH_i was measured after patients had been exposed to different levels of PEEP for a short duration. Although short-term application of high PEEP did not significantly change pH_i , it is conceivable that longer durations or higher numbers of patients would have led to more prominent reductions and statistically significant differences.

Collectively, the present findings indicate that determination of PEEP_{opt} by titration of PEEP based on improvement in oxygenation is a safe strategy, with no impairment in gastric mucosal perfusion, when cardiac output is preserved. Maintenance of cardiac output during mechanical ventilation with high PEEP may be adequate to prevent its unwanted effects

Key message

- Incremental increase in PEEP to identify the optimal value does not affect splanchnic perfusion as assessed by gastric tonometry

on organs in the splanchnic vasculature. Nonetheless, the possibility that PEEP can alter splanchnic perfusion when it is applied at high levels and for longer durations cannot be completely excluded.

Competing interests

None declared.

References

1. Guyton AC, Lindsey AW, Abernathy B, Richardson T: **Venous return at various right atrial pressure and the normal venous return curve.** *Am J Physiol* 1957, **189**:609-615.
2. Pastores SM, Katz DP, Kvetan V: **Splanchnic ischemia and gut mucosal injury in sepsis and the multiple organ dysfunction syndrome.** *Am J Gastroenterol* 1996, **91**:1697-1710.
3. Mutlu GM, Mutlu EA, Factor P: **GI complications in patients receiving mechanical ventilation.** *Chest* 2001, **119**:1222-1241.
4. Brienza N, Revelly JP, Ayuse T, Robotham JL: **Effects of PEEP on liver arterial and venous blood flows.** *Am J Respir Crit Care Med* 1995, **152**:504-510.
5. Fujita Y: **Effects of PEEP on splanchnic hemodynamics and blood volume.** *Acta Anaesthesiol Scand* 1993, **37**:427-431.
6. Arvidsson D, Almquist P, Haglund U: **Effects of positive end-expiratory pressure on splanchnic circulation and function in experimental peritonitis.** *Arch Surg* 1991, **126**:631-636.
7. Winso O, Biber B, Gustavsson B, Holm C, Milsom I, Niemand D: **Portal blood flow in man during graded positive end-expiratory pressure ventilation.** *Intensive Care Med* 1986, **12**:80-85.
8. Bonnet F, Richard C, Glaser P, Lafay M, Guesde R: **Changes in hepatic flow induced by continuous positive pressure ventilation in critically ill patients.** *Crit Care Med* 1982, **10**:703-705.
9. Kiefer P, Nunes S, Kosonen P, Takala J: **Effect of positive end-expiratory pressure on splanchnic perfusion in acute lung injury.** *Intensive Care Med* 2000, **26**:376-383.
10. Bernard GR, Argytas A, Brigham KL, Carlet J, Falke K, Hudson L, Lamy M, LeGall JR, Morris A, Spragg R, and the Consensus Committee: **The American-European Consensus Conference on ARDS. Definition, mechanism, relevant outcomes and clinical trial coordination.** *Am J Respir Crit Care Med* 1994, **149**:818-824.
11. Takala J, Parviainen I, Siloaho M, Ruokonen E, Hamalainen E: **Saline PCO_2 is an important source of error in the assessment of gastric intramucosal pH.** *Crit Care Med* 1994, **22**:1877-1879.
12. Fiddian-Green RG: **Tonometry: theory and applications.** *Intensive Care World* 1992, **9**:60-65.
13. Matuschak GM, Pinsky MR, Roger RM: **Effects of positive end-expiratory pressure on hepatic blood flow and performance.** *J Appl Physiol* 1987, **62**:1377-1383.
14. Berendes E, Lippert G, Loick HM, Brussel T: **Effects of positive end-expiratory pressure ventilation on splanchnic oxygenation in humans.** *J Cardiothorac Vasc Anesth* 1996, **10**:598-602.
15. Geiger K, Georgieff M, Lutz H: **Side effects of positive pressure ventilation on hepatic function and splanchnic circulation.** *Int J Clin Monit Comput* 1986, **3**:103-106.
16. Schlichtig R, Mehta N, Gayowski TJ: **Tissue-arterial PCO_2 difference is a better marker of ischemia than intramural pH (pHi) or arterial pH- pHi difference.** *J Crit Care* 1996, **11**:51-56.
17. Bernardin G, Lucas P, Hyvernat H, Deloffre P, Mattei M: **Influence of alveolar ventilation changes on calculated gastric intramucosal pH and gastric-arterial PCO_2 difference.** *Intensive Care Med* 1999, **25**:249-251.
18. Heard SO, Helsmoortel CM, Kent JC, Shahnarian A, Fink MP: **Gastric tonometry in healthy volunteers: effect of ranitidine on calculated intramural pH.** *Crit Care Med* 1991, **19**:271-274.

19. Kolkman JJ, Groenevelt AB, Meuswissen SG: **Effect of gastric feeding on intragastric P(CO₂) tonometry in healthy volunteers.** *J Crit Care* 1999, **14**:34-38.
20. Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, Kairalla RA, Deheinzelin D, Munoz C, Oliveira R, Takagaki TY, Carvalho CR: **Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome.** *N Engl J Med* 1998, **338**:347-354.